

BOWEL CLEANSERS

Generic	Strength	Brand	Form	Usual Dose			
HYPEROSMOTICS							
monobasic sodium phosphate	19g/118mL	Fleet Enema	OTC	enema ¹	Adults: 1 enema rectally daily; may use up to 7 days. Results usually within 1–5min; reevaluate if no relief after 7 days. Children: Use other forms.		
dibasic sodium phosphate	7g/118mL						
monobasic sodium phosphate	19g/197mL	Fleet Enema Extra					
dibasic sodium phosphate	7g/197mL						
ISOSMOTICS							
polyethylene glycol 3350	60g/L	CoLyte w. Flavor Packs	Rx	pwd for reconstitution (add 1 flavor pack before reconstitution)	Adults: After fasting for 3–4hrs, 240mL orally every 10min or 20–30mL/min by NG tube until fecal discharge is clear. Usual dose 3–4L. Only clear liqs until exam. Children: Not recommended.		
sodium chloride	1.46g/L						
potassium chloride	0.745g/L						
sodium bicarbonate	1.68g/L						
sodium sulfate	5.68g/L	GoLyteLy	Rx	pwd for reconstitution			
polyethylene glycol 3350	227.1g/4L						
sodium chloride	5.53g/4L						
potassium chloride	2.82g/4L						
sodium bicarbonate	6.36g/4L	TriLyte w. Flavor Packs	Rx	pwd for reconstitution (add 1 flavor pack before reconstitution)	Adults: May consume water or clear liquids during and after completion of bowel prep up until 2hrs before procedure. Drink 240mL every 10min or 20–30mL/min by NG tube until rectal effluent is clear or 1 gal consumed. Children: Not established.		
sodium sulfate	21.5g/4L						
polyethylene glycol 3350	420g/4L						
sodium chloride	11.2g/4L						
potassium chloride	1.48g/4L	MoviPrep	Rx	pwd for reconstitution ²		Adults: After fasting for 3–4 hrs, 240mL orally every 10min or 20–30mL/min by NG tube until rectal discharge is clear or 4L consumed. Children: <6mos: not recommended. ≥6mos: No solid food for at least 2hrs, then 25mL/kg/hr orally or by NG tube until rectal discharge is clear or 4L consumed. Both: May interrupt if severe GI disturbances occur. Only clear liq until exam.	
sodium bicarbonate	5.72g/4L						
polyethylene glycol 3350	200g/2L						
sodium sulfate	15g/2L						
sodium chloride	5.38g/2L	NuLyteLy	Rx	pwd for reconstitution			
potassium chloride	2.03g/2L						
ascorbic acid	9.4g/2L						
sodium ascorbate	11.8g/2L						
polyethylene glycol 3350	420g/4L	OsmoPrep	Rx	tabs	Adults: 240mL orally every 10min or 20–30mL/min by NG tube until rectal effluent is clear or 4L consumed. Children: <6mos: not recommended. ≥6mos: 25mL/kg/hr orally or by NG tube until rectal effluent is clear or 4L consumed. Both: May consume water or clear liq during and after completion of bowel prep up until 2hrs before procedure.		
sodium bicarbonate	5.72g/4L						
sodium chloride	11.2g/4L						
potassium chloride	1.4g/4L						
sodium phosphate (as monobasic monohydrate and dibasic anhydrous)	1.5g	Plenvu	Rx	pwd for reconstitution		Adults: After fasting for 3–4 hrs, 240mL orally every 10min or 20–30mL/min by NG tube until rectal discharge is clear or 4L consumed. Children: <6mos: not recommended. ≥6mos: 25mL/kg/hr orally or by NG tube until rectal discharge is clear or 4L consumed. Both: May consume water or clear liq during and after completion of bowel prep up until 2hrs before procedure.	
polyethylene glycol 3350	140g/32oz						
sodium ascorbate	48.11g/32oz						
sodium sulfate	9g/32oz						
ascorbic acid	7.54g/32oz	Suprep	Rx	soln for dilution			Adults: Evening before colonoscopy: 4 tabs with 8oz of clear liq every 15mins for a total of 20 tabs; repeat on day of colonoscopy 3–5hrs before procedure for a total of 12 tabs. Wait 7 days before retreating. <18yrs: Not recommended.
sodium chloride	5.2g/32oz						
potassium chloride	2.2g/32oz						
sodium sulfate	17.5g/6oz						
potassium sulfate	3.13g/6oz	OsmoPrep	Rx	tabs	Adults: Two-day regimen: start the evening before colonoscopy, drink 480mL over 30mins, follow with 480mL of clear liquids over next 30mins; in the AM, drink 480mL over 30mins, then drink 480mL of clear liquids over next 30mins. Consume additional water or clear liquids up to 2hrs before colonoscopy. One-day regimen: start the morning of colonoscopy, drink 480mL over 30mins, follow with 480mL of clear liquids over next 30mins. At minimum of 2hrs after first dose, drink 480mL over 30mins, follow with 480mL of clear liquids over next 30mins. Consume additional water or clear liquids up to 2hrs before colonoscopy. Children: Not established.		
magnesium sulfate	1.6g/6oz						
polyethylene glycol 3350	140g/32oz						
sodium ascorbate	48.11g/32oz						
sodium sulfate	9g/32oz	Suprep	Rx	soln for dilution		Adults: Start the evening before colonoscopy. Dilute 1 bottle to 16oz with water and drink entire amount. Drink 32oz of water over the next hour. Next morning: repeat using the second bottle. Complete prep at least 2hrs before colonoscopy. Children: Not established.	
ascorbic acid	7.54g/32oz						
sodium chloride	5.2g/32oz						
potassium chloride	2.2g/32oz						
sodium sulfate	17.5g/6oz	Suprep	Rx	soln for dilution			Adults: Start the evening before colonoscopy. Dilute 1 bottle to 16oz with water and drink entire amount. Drink 32oz of water over the next hour. Next morning: repeat using the second bottle. Complete prep at least 2hrs before colonoscopy. Children: Not established.
potassium sulfate	3.13g/6oz						
magnesium sulfate	1.6g/6oz						
polyethylene glycol 3350	140g/32oz						
sodium ascorbate	48.11g/32oz	Plenvu	Rx	pwd for reconstitution	Adults: Two-day regimen: start the evening before colonoscopy, drink 480mL over 30mins, follow with 480mL of clear liquids over next 30mins; in the AM, drink 480mL over 30mins, then drink 480mL of clear liquids over next 30mins. Consume additional water or clear liquids up to 2hrs before colonoscopy. One-day regimen: start the morning of colonoscopy, drink 480mL over 30mins, follow with 480mL of clear liquids over next 30mins. At minimum of 2hrs after first dose, drink 480mL over 30mins, follow with 480mL of clear liquids over next 30mins. Consume additional water or clear liquids up to 2hrs before colonoscopy. Children: Not established.		
sodium sulfate	9g/32oz						
ascorbic acid	7.54g/32oz						
sodium chloride	5.2g/32oz						
potassium chloride	2.2g/32oz	Suprep	Rx	soln for dilution		Adults: Start the evening before colonoscopy. Dilute 1 bottle to 16oz with water and drink entire amount. Drink 32oz of water over the next hour. Next morning: repeat using the second bottle. Complete prep at least 2hrs before colonoscopy. Children: Not established.	
sodium sulfate	17.5g/6oz						
potassium sulfate	3.13g/6oz						
magnesium sulfate	1.6g/6oz						
STIMULANTS							
bisacodyl	10mg/30mL	Fleet Bisacodyl Enema	OTC	enema ¹			Adults: 1 enema rectally daily. Results usually within 5–20min; reevaluate if ineffective. Children: Not recommended.
STIMULANT + OSMOTIC LAXATIVE							
sodium picosulfate	10mg	Clenpiq	Rx	oral soln			Adults and Children: <9yrs: not established. ≥9yrs: Split-Dose regimen (preferred): Take 1st dose evening before colonoscopy followed by five 8oz cups of clear liquids within 5hrs and before bed. Take 2nd dose, the next day approx. 5hrs before colonoscopy followed by at least three 8oz cups of clear liquids, ≥2hrs before procedure. Day-Before regimen (alternative): Take 1st dose in the afternoon or early evening before colonoscopy followed by five 8oz cups of clear liquids within 5hrs and before the next dose. Take 2nd dose approx. 6hrs later in the evening, the night before colonoscopy followed by three 8oz cups of clear liquids within 5hrs and before bed.
magnesium oxide	3.5g						
anhydrous citric acid	12g						
sodium picosulfate	10mg/pkt	Prepopik	Rx	pwd for reconstitution			
magnesium oxide	3.5g/pkt						
anhydrous citric acid	12g/pkt						

NOTES

Key: e-c = enteric-coated; del-rel = delayed-release; pwd = powder; supp = suppository

¹latex free, contains sodium 4.4g/enema

²contains phenylalanine 131mg/treatment

Not an inclusive list of medications and/or official indications. Please see drug monograph at www.eMPR.com and/or contact company for full drug labeling.

(Rev. 8/2019)