

PARKINSONISM TREATMENTS (Part 1 of 2)

Generic	Brand	Form	Strength	Adult Dose
ANTICHOLINERGICS				
benztropine	—	scored tabs	0.5mg, 1mg, 2mg	Initially 0.5–1mg at bedtime. May increase by 0.5mg at 5–6 day intervals; max 6mg daily.
	Cogentin	amps	1mg/mL	
hyoscyamine sulfate	NuLev	chewable melt tabs	0.125mg	1–2 tabs every 4hrs or as needed; max 12 tabs/day
trihexyphenidyl	—	scored tabs	2mg, 5mg	Give in 3–4 divided doses. 1mg on Day 1, may increase by 2mg every 3–5 days; usual max 6–15mg/day. <i>Concomitant L-dopa</i> : 3–6mg/day and reduce L-dopa dose.
		susp	2mg/5mL	
CATECHOL O-METHYL TRANSFERASE (COMT) INHIBITORS				
entacapone	Comtan	tabs	200mg	200mg with each dose of L-dopa/carbidopa, up to 8 times daily
tolcapone	Tasmar	tabs	100mg	100mg three times daily; may cautiously increase to 200mg three times daily
CHOLINESTERASE INHIBITORS				
rivastigmine	Exelon	caps	1.5mg, 3mg, 4.5mg, 6mg	Initially 1.5mg twice daily (AM & PM); if tolerated, may increase by 1.5mg twice daily at intervals of at least 4 weeks. Usual range: 3–12mg/day; max 12mg/day.
		patches	4.6mg/24hrs, 9.5mg/24hrs, 13.3mg/24hrs	Initially apply one 4.6mg/24hrs patch once daily; if tolerated, may increase to 9.5mg/24hrs patch after 4 weeks at previous dose; can further increase to max 13.3mg/24hrs dose.
DOPA-DECARBOXYLASE INHIBITOR				
carbidopa	Lodosyn	tabs	25mg	<i>Concomitant Sinemet 10-100</i> : 25mg with first dose of Sinemet each day; additional 12.5mg or 25mg doses may be given with each dose of Sinemet. <i>Concomitant Sinemet 25-100 or 25-250</i> : 25mg with any dose of Sinemet as required for optimum therapeutic response. Max total carbidopa 200mg/day.
DOPA-DECARBOXYLASE INHIBITOR + DOPAMINE PRECURSOR				
carbidopa/levodopa*	Duopa	enteral susp	4.63mg/20mg per mL	Day 1: calculate and administer initial daily (Morning Dose + Continuous Dose); titrate subsequent doses based on response. Max daily dose: 2000mg of levodopa (1 cassette) over 16hrs. See full labeling.
	Rytary	ext-rel caps	23.75mg/95mg, 36.25mg/145mg, 48.75mg/195mg, 61.25mg/245mg	<i>Levodopa-naive</i> : Initially 23.75mg/95mg 3 times daily for the first 3 days; may increase to 36.25mg/145mg 3 times daily on the 4th day; up to max 97.5mg/390mg 3 times daily. May increase to max 5 times daily if more frequent dosing needed and tolerated. Max daily dose: 612.5mg/2450mg.
	—	ODT	10mg/100mg, 25mg/100mg, 25mg/250mg	Initially one 25mg/100mg tab 3 times daily, or one 10mg/100mg tab 3–4 times daily; increase every 1–2 days up to 2 tabs (of either 25/100 or 10/100) 4 times daily. <i>Patients taking L-dopa > 1500mg/day</i> : Initially one 25mg/250mg tab 3–4 times daily; max carbidopa 200mg/day. <i>For ODT</i> : Discontinue levodopa at least 12hrs before.
	Sinemet	tabs	10mg/100mg, 25mg/100mg, 25mg/250mg	
	Sinemet CR	sust-rel tabs	25mg/100mg, 50mg/200mg+	<i>Not receiving L-dopa</i> : Initially one 50mg/200mg tab twice daily, at intervals of at least 6hrs. Allow 3 days between dosage adjustments. If given at intervals <4hrs and/or divided doses not equal: give smaller doses at end of day. May add immediate-release Sinemet 25-100 or 10-100 tabs in advanced disease.
DOPAMINE AGONISTS				
amantadine	—	tabs	100mg	<i>Monotherapy</i> : 100mg twice daily; may increase after 1–2wks by 100mg daily. <i>Serious associated illness or high doses of other antiparkinson drugs</i> : 100mg once daily, may increase after 1 to several weeks to 100mg twice daily; max 400mg/day in divided doses. <i>Renal dysfunction</i> : Reduce dose; see full labeling.
		susp	50mg/5mL	
apomorphine	Apokyn	cartridges	10mg/mL	Supervise 1st dose (monitor BP); prescribe for outpatient use at a dose at least 0.1mL less than tolerated test dose; usual range 0.2mL to 0.6mL; max 0.6mL/episode and one dose/episode; usual max 5 doses/day (2mL/day). See full labeling.

(continued)

PARKINSONISM TREATMENTS (Part 2 of 2)

Generic	Brand	Form	Strength	Adult Dose
DOPAMINE AGONISTS (continued)				
bromocriptine	Parlodel	caps	5mg	Initially 1.25mg twice daily. May increase every 2–4wks by 2.5mg/day; max 100mg/day.
		scored tabs	2.5mg	
pramipexole*	Mirapex	tabs	0.125mg, 0.25mg+, 0.5mg+, 0.75mg, 1mg+, 1.5mg+	0.125mg three times daily. May increase gradually at intervals of 5–7 days up to max 1.5mg three times daily. <i>Renal impairment (CrCl 30–50mL/min):</i> 0.125mg three times daily. <i>CrCl 15–<30mL/min:</i> 0.125mg once daily; max 1.5mg once daily. <i>CrCl <15mL/min, hemodialysis: not recommended.</i>
	Mirapex ER	ext-rel tabs	0.375mg, 0.75mg, 1.5mg, 2.25mg, 3mg, 3.75mg, 4.5mg	0.375mg once daily; may increase gradually at intervals of 5–7 days, first to 0.75mg/day, then by 0.75mg increments up to max 4.5mg/day. <i>Renal impairment (CrCl 30–50mL/min):</i> give every other day; reevaluate before increasing to daily dosing after 1 week and before titrating by 0.375mg increments up to 2.25mg/day. <i>CrCl <30mL/min, hemodialysis: not recommended.</i>
ropinirole*	Requip	tabs	0.25mg, 0.5mg, 1mg, 2mg, 3mg, 4mg, 5mg	0.25mg 3 times daily, then increase by 0.25mg 3 times daily at 1wk intervals to 1mg 3 times daily to 4th week. May increase by 1.5mg/day at 1wk intervals up to 9mg/day, then by up to 3mg/day at 1wk intervals to max 24mg/day.
	Requip XL	ext-rel tabs	2mg, 4mg, 6mg, 8mg, 12mg	2mg once daily for 1–2wks, then increase by 2mg/day at ≥1wk intervals up to max 24mg/day (for advanced disease: usually up to max 8mg/day; early disease: usually up to max 12mg/day).
rotigotine	Neupro	patches	1mg/24hrs, 2mg/24hrs, 3mg/24hrs, 4mg/24hrs, 6mg/24hrs, 8mg/24hrs	<i>Early-stage:</i> Initially 2mg/24hrs patch once daily; may increase weekly by 2mg/24hrs if needed; max 6mg/24hrs once daily. <i>Advanced-stage:</i> Initially 4mg/24hrs patch once daily; may increase weekly by 2mg/24hrs if needed; max 8mg/24hrs once daily.
DOPA-DECARBOXYLASE INHIBITOR + DOPAMINE PRECURSOR + COMT INHIBITORS				
carbidopa/levodopa/entacapone	Stalevo	tabs	12.5mg/50mg/200mg, 18.75mg/75mg/200mg, 25mg/100mg/200mg, 31.25mg/125mg/200mg, 50mg/200mg/200mg, 37.5mg/150mg/200mg	Max 1 tab per dosing interval. <i>Previously on carbidopa/levodopa and entacapone:</i> Substitute on a mg/mg basis. Stalevo 50, 75, 100, 125, 150: max 8 tabs/day; Stalevo 200: max 6 tabs/day.
DOPAMINE PRECURSOR				
levodopa	Inbrija	caps	42mg	Inhale contents of 2 caps (84mg) as needed, up to 5 times daily. Max dose per OFF period: 84mg; max daily dose: 420mg.
MONOAMINE OXIDASE-B INHIBITORS				
rasagiline	Azilect	tabs	0.5mg, 1mg	<i>Monotherapy or adjunct w/o levodopa:</i> 1mg once daily. <i>Concomitant levodopa with/without other PD drugs (eg, dopamine agonist, amantadine, anticholinergics):</i> Initially 0.5mg once daily; may increase to 1mg once daily (consider reducing levodopa dose based on response). <i>Mild hepatic impairment (Child-Pugh score 5–6) or concomitant CYP1A2 inhibitors:</i> 0.5mg once daily.
safinamide	Xadago	tabs	50mg, 100mg	<i>Adjunct to levodopa/carbidopa:</i> Initially 50mg once daily; may increase to 100mg once daily after 2wks as tolerated. <i>Moderate hepatic impairment (Child-Pugh B):</i> max 50mg once daily.
selegiline	—	caps	5mg	5mg at breakfast and at lunch; max 10mg/day. After 2–3 days, L-dopa/carbidopa dosage may be reduced by 10–30%.
	Zelapar	ODT	1.25mg	1.25mg once in the AM for at least 6wks; if needed, may increase to max 2.5mg once daily if tolerated.

NOTES

Key: susp = suspension; ODT = orally disintegrating tablets; sust-rel tabs = sustained release tablets; ext-rel tabs = extended release tablets; soln = solution; amps = ampules; + = scored tablets.

*First line treatment for Parkinson's disease.

Not an inclusive list of medications and/or official indications. Please see drug monograph at www.eMPR.com and/or contact company for full drug labeling.