

CHRONIC OBSTRUCTIVE PULMONARY DISEASE TREATMENTS

Risk Factors: genetic factors, exposure to particles (tobacco smoke, occupational dusts, outdoor air pollution), oxidative stress, respiratory infections, nutrition, comorbidities.

Classification

- Stage 1: Mild – FEV₁/FVC < 0.70; FEV₁ ≥ 80% predicted.
- Stage 2: Moderate – FEV₁/FVC < 0.70; 50% ≤ FEV₁ < 80% predicted.
- Stage 3: Severe – FEV₁/FVC < 0.70; 30% ≤ FEV₁ < 50% predicted.
- Stage 4: Very Severe – FEV₁/FVC < 0.70; FEV₁ < 30% predicted or FEV₁ < 50% predicted + CRF.

Step-wise treatment: Visit the Global Initiative for Chronic Obstructive Lung Disease (GOLD) website at www.goldcopd.org for information about use of medications at various stages of COPD.

Generic	Brand	Form	Usual Dosage
---------	-------	------	--------------

BRONCHODILATORS

Long-Acting Beta₂-Agonists (LABAs)

arformoterol	Brovana ¹	soln	Adults: Inhale 15mcg twice daily (AM & PM) by nebulization (max: 30mcg/day). Use standard jet nebulizer with air compressor (see full labeling). Children: Not established.
formoterol	Perforomist ¹	soln	Adults: Inhale one 20mcg vial twice daily (AM & PM) by nebulization. Use standard jet nebulizer and air compressor. Children: Not established.
olodaterol	Striverdi Respimat ¹	MDI	Adults: 2 inh once daily; max 2 inh/24hrs Children: Not established.
salmeterol	Serevent Diskus ²	DPI	Adults: 1 inh (50mcg) twice daily (approx. 12hrs apart) Children: Not established.

Short-Acting Anticholinergics

ipratropium bromide	— ¹	soln	Adults: 500mcg by oral nebulization 3–4 times daily every 6–8hrs Children: Not established.
	Atrovent HFA ¹	MDI	Adults: 2 inh 4 times daily (max: 12 inh/day) Children: Not established.

Long-Acting Anticholinergics

acclidinium bromide	Tudorza Pressair ¹	DPI	Adults: 1 inh (400mcg) twice daily (approx. 12hrs apart) Children: Not established.
glycopyrrolate	Lonhala Magnair ¹	soln	Adults: 1 oral inh of one 25mcg vial twice daily (AM & PM), using Magnair device Children: Not established.
revefenacin	Yupelri ¹	soln	Adults: 1 oral inh of one 175mcg vial once daily using a mouthpiece. Use standard jet nebulizer with air compressor. Children: Not established.
tiotropium bromide	Spiriva HandiHaler ¹	caps	Adults: 2 inh of one 18mcg caps once daily, using HandiHaler device. Max: 2 inh/24hrs. Children: Not established.
	Spiriva Respimat	MDI	Adults: 2 inh of 2.5mcg/actuation (5mcg) once daily; max: 2 inh/24hrs Children: Not established.
umeclidinium	Incruse Ellipta ¹	DPI	Adults: 1 inh every 24hrs Children: Not established.

Anticholinergic + Beta₂-Agonist

ipratropium bromide + albuterol	— ¹	soln	Adults: ≥18yrs: 1 vial 4–6 times daily via nebulizer Children: <18yrs: Not established.
	Combivent Respimat ¹	MDI	Adults: 1 inh 4 times daily (max: 6 inh/day) Children: Not established.

Anticholinergic + Long-Acting Beta₂-Agonist (LABA)

acclidinium bromide/ formoterol	Duaklir Pressair ^{1,2}	dry pwd	Adults: 1 inh twice daily (AM & PM); max 1 inh twice daily Children: Not established.
glycopyrrolate + formoterol	Bevespi Aerosphere ^{1,2}	MDI	Adults: 2 inh twice daily (AM & PM); max 2 inh twice daily Children: Not established.
tiotropium + olodaterol	Stiolto Respimat ¹	MDI	Adults: 2 inh once daily (max: 2 inh/24hrs) Children: Not established.
umeclidinium + vilanterol	Anoro Ellipta ^{1,2}	DPI	Adults: 1 inh once daily Children: Not established.

CORTICOSTEROIDS

Corticosteroid + Long-Acting Beta₂-Agonist (LABA)

budesonide + formoterol	Symbicort 160/4.5 ²	MDI	Adults: 2 inh of 160/4.5mcg twice daily (approx. 12hrs apart) Children: Not indicated.
fluticasone + salmeterol	Advair 250/50 Diskus ^{2,3}	DPI	Adults: 1 inh of 250/50mcg twice daily (approx. 12hrs apart) Children: Not established.
	Wixela Inhub ²	DPI	Adults: 1 inh of 250/50mcg twice daily (approx. 12hrs apart) Children: Not established.
fluticasone + vilanterol	Breo Ellipta ²	DPI	Adults: 1 inh of 100/25mcg once daily (max) Children: ≤17yrs: Not established.

Corticosteroid + Anticholinergic + Long-Acting Beta₂-Agonist (LABA)

budesonide + glycopyrrolate + formoterol	Breztri Aerosphere ^{1,2}	MDI	Adults: 2 inh twice daily (AM & PM); max 2 inh twice daily Children: Not established.
fluticasone + umeclidinium + vilanterol	Telegly Ellipta ²	DPI	Adults: 1 inh of 100/62.5/25mcg once daily (max) Children: Not established.

OTHER

PDE4-Inhibitor

roflumilast	Daliresp ^{1,2}	tabs	Adults: 500mcg once daily. Initiating 250mcg daily for 4wks, then increasing to 500mcg may reduce the rate of discontinuation in some patients. Children: Not established.
-------------	--------------------------------	------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

NOTES

Key: CRF = chronic respiratory failure; DPI = dry powder inhaler; FEV₁ = forced expiratory volume in one second; FVC = forced vital capacity; MDI = metered dose inhaler

¹Indicated only for COPD. ²Not indicated for the relief of acute bronchospasm. ³Only Advair 250/50 Diskus twice daily is approved for maintenance treatment of COPD because an efficacy advantage of the higher strength Advair 500/50 over Advair 250/50 has not been demonstrated. Other strengths and formulations of Advair are available.

Not an inclusive list of medications and/or official indications. Please see drug monograph at www.eMPR.com and/or contact company for full drug labeling.

(Rev. 7/2021)